

Avicenniaceae Verbena family

Jorge A. Jimenez and Ariel E. Lugo

HABITAT

Native Range

Avicennia germinans (L.) L., the black mangrove, occurs in most of the mangrove forests of the American coastal areas (fig. 1). It is found throughout the coast of the Gulf of Mexico, and from northern Florida (29°53' N) to Espiritu Santo, Brazil (approximately 23° S). On the Pacific coasts of North and South America, it grows from Punta de Lobos, Mexico (30°15' N), to south of Punta Malpelo, Peru (3°40'S) (11, 64).

Black mangrove grows in tidal areas with salty or brackish waters. It grows well in basin forests over a broad range of soil salinities (23, 30). The best structural development is attained in tropical wet riverine forests (23). Black mangrove also grows in the less flushed inner portions of fringe and riverine forests. In these areas it shows a distinct preference for slightly higher drier soils (55). At the geographical limit of mangrove distribution, black mangroves occupy the forest fringe (33).

Climate

Black mangroves grow over a wide variety of climates. They grow in tropical and subtropical dry, moist, and wet life zones, with a broad range of precipitation regimes (from 800 to 7,000 mm per year).

The species is sensitive to frost, but of all the mangrove species, it is considered the most tolerant to low temperatures (12). In its northern limits of distribution, black mangroves are killed by freezing temperatures -3 to -11°C (33, 64). In these areas, tree heights are shorter relative to trees growing at lower latitudes. Black mangrove is the dominant species on sites with arid climates, where soil salinities exceed 40 parts per thousand (13).

Soils and Topography

Black mangrove distribution is highly influenced by changes in microtopography and the consequent changes in soil flooding and salinity (10, 14).

The species is usually found in low areas, inland from the fringe of the mangrove forest. However, it also grows inland, on slightly elevated areas, where tidal flooding

UNESCO fellow and Project Leader, respectively, Institute of Tropical Forestry, Southern Forest Experiment Station, USDA Forest Service, Rio Piedras, Puerto Rico.

is less frequent. Here soils are covered by a few centimeters of continuously standing water or are inundated only a few times per year (213 to 432 tides per year, 10; 152 to 158 tides per year, 58).

Black mangrove grows on sandy, silty, or clay soils. It is found on heavily oxidized clays or on soils with significant concentrations of pyrite (2). Black mangrove soils have organic matter contents of 2 to 25 percent (23), but values may be as high as 58 percent (34, 58); nitrogen content is low—around 0.4 percent (20).

The species can grow on soils where salinities range from 0 to 100 parts per thousand (51). Under high soil salinities, structural development is suppressed (13). Leaves excrete salt through specialized glands and may be salt covered, thus contributing to salty through-fall (31).

Associated Forest Cover

Black mangrove may be found in pure stands or in close association with other mangrove species within its range. It grows with *Rhizophora mangle* L., *R. harrisonii* Leechm., *R. racemosa* G.F.W. Meyer, *Avicennia*



Figure 1.—Distribution of *Avicennia germinans* in the New World.

tonduzzii Moldenke, *A. schaueriana* Stapf & Leechm., *A. bicolor* Standl., *Laguncularia racemosa* Gaertn., and *Pelliciera rhizophorae* Tr. & Pl.

In basin forests, where soil salinities are around 30 to 40 parts per thousand, black mangrove grows with white mangrove (*L. racemosa*); if soil salinities are higher than 50 parts per thousand, the black mangrove will be dominant (14). In areas of low soil salinity, black mangrove may be associated with *Pterocarpus officinalis* Jacq., *Mora oleifera* (Triana) Duke, *Conocarpus erecta* L., and the fern *Acrostichum aureum* Troll, Sloane, Hooker.

LIFE HISTORY

Reproduction and Early Growth

Flowering and Fruiting.—Flowers are found in axillary and terminal inflorescences, with 1 to 15 pairs of flowers per spike. They are small (1 to 2 cm wide), sessile, and with imbricate bracts. The corolla has four lobes. Petals are yellow or cream to white, generally with a yellowish throat (10, 38). The fragrant insect-pollinated flower has a two-celled ovary; each loculus contains two locules (39). Flowering is sporadic throughout the year, even though distinct peaks can be observed (36, 45). Sexual maturity is reached when plants are 2 to 3 m tall (26).

The species is considered viviparous because germination occurs while the embryo is still enclosed in the fruit. A distinctive seedling develops before the fruit falls from the parent tree. The expansion of the cotyledons and the development of plumular leaves is evident (54).

Three of the four ovules abort, producing a one-seeded fruit. The fruit is oblong or elliptic with an approximate weight of 1.1 g and an average length of 1.8 cm (6). Trees have been reported to produce over 300 fruits per year (10).

Fallen seedlings float and are transported by tidal currents. The propagule sheds its pericarp and produces roots within 3 weeks of dispersal (45). Seedlings of *Avicennia* can become waterlogged (15). Establishment is limited to areas above water level at low tide (64). Water turbulence inhibits root development, and high water temperatures (39° to 40°C) that last over 48 hours are lethal to the seedlings (37).

Densities of 0.07 seedlings/m² (34, 58) and establishment rates of 0.06 seedlings/m²·yr (4) have been reported. Cohorts of black mangrove seedlings exhibit up to 30 percent survival (17, 46).

Vegetative Reproduction.—Black mangrove coppices fairly well, provided the stump is not submerged (36). Air layering techniques applied to this species have had little success; only 6 percent of the trees produced roots and rootlets (6).

Sapling and Pole Stage to Maturity

Growth and Yield.—Black mangrove growth is characterized by continuous growth of the axes (i.e. Attimis's

model, 19). Branching is diffuse with slightly slanting secondary axes (26). Growth rings are produced, and the number of rings and the average diameter of the section appear to be related. However, the number of rings are not related to age (18). Diameter-age measurements for black mangrove individuals in the Caribbean show average diameter at breast height (d.b.h.) of 8.6, 12.2, and 19.1 cm for trees 10, 20, and 50 years old, respectively (43).

Planting of wildlings (0.6 m tall) has given poor results (21). *Avicennia* spp. in the Indo-Pacific region are planted by planting seedlings during periods of low tide (56).

Plants 0.5 to 1.5 m tall, transplanted with a rootball 1.5 times wider than tree height, had good recovery (44). Pruned saplings grew 3.9 times faster than unpruned controls (44). Tree stature varies enormously, with some mature (reproductively active) individuals only 20 cm tall and others growing to a height of 36 m with a d.b.h. of 1.8 m (fig. 2). Climatic and edaphic conditions are responsible for this variability (13, 23). In the Caribbean, black mangrove forests generally do not exceed 15 m, even though on the wet Central American coast, individual trees more than 30 m in height can be found. Monospecific black mangrove basin forests in south Florida have basal areas of 12.5 to 21.2 m²/ha, and tree density ranges from 2,467 to 6,511 trees/ha (34, 58).

In riverine forests in Venezuela, parcels dominated by black mangroves had wood volumes of up to 350 m³/ha (35). In Puerto Rico, periodic annual diameter increment was 0.42, 0.46, and 0.51 cm per year in three replicate stands over a 37-year period after clearcutting (62). Periodic basal area growth (table 1) decreased with age. These results are for black mangrove growing in stands dominated by white mangrove (*L. racemosa*).

In Venezuelan riverine forests, 20-by 300-m parcels rotated every 30 years are used as management units (35). *Avicennia* spp. in India have been used with a clear-felling system and 15 to 25 year rotations. Here, 50 to 60 trees/ha are left as seed trees (12).

Net primary production rates ranging from 0.94 to 2.7 gC/m²·day have been measured in black mangrove forests growing in water with a low chlorine content (5 to 16 parts per thousand) (7). Gross primary productivity (averaging 8.62 gC/m²·day) (31) increases with increasing chlorinity between the range of 5 to 16 parts per thousand (7). Wood production in a disturbed forest was 1.74 g/m²·day (31). The complexity index (22) of black mangrove stands in south Florida decreased from 40 to 3.4 when soil salinities increased from 40 to 65 parts per thousand (34, 58). Water loss via transpiration from black mangroves was 2.53 mm per day relative to 4.19 mm per day for adjacent red mangroves (31).

Rooting Habit.—Black mangrove is characterized by a shallow underground root system with thin sinker roots and negatively geotropic pneumatophores that develop from lateral horizontal roots (24). Soft, slender roots occasionally emerge from the trunk of old trees (39, 50). Pneumatophores are responsible for gas ex-




Figure 2.—A mature *Avicennia germinans* tree.

change processes (32, 49). An average density of 672 pneumatophores/m² has been measured in basin forests of south Florida (4, 58). However, the actual number of pneumatophores is highly variable (2). Pneumatophore height increases with increasing water depth (29).

Root biomass in *Avicennia* species may account for up to 65 percent of the total tree biomass (16). Fibrous roots comprise approximately 50 percent, while pneumatophores and lateral roots each comprise approximately 25 percent of the total root biomass (16). Fast root production rates allow the species to become established on rapidly accreting shorelines and adjust quickly to rapid sedimentation (64).

Reaction to Competition.—Black mangrove is considered intolerant of shade and is unable to regenerate in even moderate shade (61, 45). In areas where there is constant flushing, black mangrove is dominated by other mangrove species. In the inner parts of the forest under low soil salinity conditions, brackish water species such as *Pterocarpus officinalis* and *Mora oleifera* out-compete black mangrove (12).

Damaging Agents.—The wood borer *Sphaeroma terebrans* Bate has been found in exposed roots of black mangrove trees (48). The fungi *Alternaria alternata* and *Phytophthora* spp. have been reported as the cause

Table 1.—Basal area growth in three Puerto Rican mangrove stands clearcut in 1937. *Avicennia* comprised between 2 to 6 percent of the stems in 1938 and between 20 to 30 percent in 1975.¹

Time interval	<i>Laguncularia racemosa</i>	<i>Avicennia germinans</i>	Total
years	m ² /ha. yr		
1938–1945	0.18
	1.20
	1.48
1945–1949	2.8	0.44	3.24
	2.17	0.37	2.54
	0.92	0.29	1.21
1951–1955	0.77	0.58	1.35
	0.71	0.31	1.02
	0.63	0.48	1.11
1955–1975	0.67	0.08	0.75
	0.94	0.12	1.06
	0.71	–0.04	0.67

¹ Data adapted from Weaver 1979 (62).

of defoliation and death in Australian *Avicennia* spp. (9, 41). Leaf infestation and defoliation by the scale *Icerya seychellarum* Westw. and the caterpillar *Cleora injectaria* Walker have been observed in *Avicennia* spp. of the Indo-Pacific (40, 42). It is common to see a high intensity of leaf miner activity in black mangrove trees.

Black mangrove is highly susceptible to changes in hydrological patterns. Drought or flooding can cause extensive mortality (3). The species is easily wind thrown and is killed when exposed to low concentrations of auxin-type herbicides (57, 60). The wood is susceptible to attack by dry-wood termites (28). Others report the wood to be termite resistant but heavily damaged by fungus and marine borers (1, 52).

SPECIAL USES

Black mangrove is used as a source of firewood and charcoal in many coastal areas of tropical America. It is also used for cheap beams, door frames, boat construction, piers, wharf posts, telegraph and power poles, and crossties (12, 28, 59). The wood responds adequately to impregnation treatment with creosote (1) and with copperized chromated borate and copperized chromated arsenate preservatives (25).

The wood has a specific gravity of 0.8 to 1.0 and a radial and tangential shrinkage of 7.1 and 10.2 percent respectively. Its hardness and modulus of rupture are classified as moderate (1, 53). Black mangrove wood is coarse and of uneven grain. The heartwood contains a lapachol compound that gives it a yellowish coloration (36). Its wood parenchyma is sparingly paratracheal and wood fibers are usually thick walled (47).

Black mangrove wood has been reported as suitable for fiberboard and paper pulp (53). It can be pulped successfully by the soda process; however, it must be mixed with other types of wood because of its short

fiber (47). Alpha cellulose contents are generally about 69 percent. The wood responds adequately to air drying.

Black mangrove wood has been reported as difficult to work (47), but others (1) have described it as adequate for saw milling. Little abrasion was observed in steel saws used to cut black mangrove wood at 22 m/s and a 40° cutting angle (1).

The bark of black mangrove is used as a source of tannins even though soluble tannin contents are only 5.5 to 12.7 percent, on a dry weight basis. Infusions from the bark are used as an astringent. The resin is used for treatment of ulcers, hemorrhoids, diarrhea, and tumors (12). Species of *Avicennia* are favored by beekeepers because they contribute to yields of good quality honey (8). Sprouting seeds are edible when cooked but poisonous raw (28). Salt for cooking or eating purposes can be obtained from salt-coated foliage (28).

Soil under *Avicennia* forests have been reported as suitable for land reclamation in Africa. The lower sulphides and organic matter contents of these soils prevent the drastic reduction of pH observed after drainage of soils in other mangrove forests (20).

Black mangrove is considered a soil stabilizer. In areas near the mean high tide level, black mangrove is believed to reduce current velocities and promote sedimentation (5, 63). Along rapidly prograding coasts, black mangroves are responsible for holding the deposited material (2, 63).

Black mangrove can also be a fast leaf litter producer. Litter-fall rates in monospecific mangrove forests show marked seasonality with peak production during the rainy season. Maximum rates fluctuate from 2.5 to 3.3 g/m²·day, while minimum rates vary from 0.4 to 0.5 g/m²·day (34, 58). Annual rates of litter fall in monospecific black mangrove forests of south Florida average 4.5 to 4.7 t/ha (34, 58). The decomposition of this material results in a highly nutritious detritus that supports secondary production in adjacent estuarine waters (58). Black mangrove leaves leach organic matter readily and have low carbon to nitrogen ratios resulting in fast decomposition rates (34, 58). These conditions enhance the development of a complex food web that includes many commercially important species of fish, mollusks, and crustaceans (58).

GENETICS

Population Differences

The genus *Avicennia* has been reviewed by Moldenke (38, 39). This author separated the genus from the Verbenaceae, creating a new family, Avicenniaceae. Four species of *Avicennia* have been reported in tropical America: *A. germinans* (L.) L., *A. bicolor* Standl., *A. tonduzzii* Moldenke, and *A. schaueriana* Stapf & Leecm.. Little (27) maintains *A. nitida* in place of *A. germinans* (L.) Stearn. Many authors consider *A. africana* P. Beauv., a dominant species along the coast of West Africa, as being the same species as *A. germinans* (L.) L.

The species is highly polymorphic, and large variation exists among populations growing under different conditions. Leaf size is one of the most variable parameters; some of this variation is due to soil salinity.

LITERATURE CITED

1. Arroyo, Joel P. 1970. Propriedades y usos posibles de los mangles de la region del rio San Juan en la reserva forestal de Guarapiche. Boletin del Instituto Latino-Americano de Investigacion y Capacitacion 33/34:53-76.
2. Augustinus, P.G.E.F. and S. Slager. 1971. Soil formation in swamp soils of the coastal fringe of Surinam. Geoderma 6:203-211.
3. Bacon, Peter R. 1970. The ecology of Caroni swamp, Trinidad. Special Publication. Central Statistical Office Trinidad. 68 p.
4. Banner, A. 1977. Revegetation and maturation of restored shorelines in Indian River. In Proceedings, fourth annual conference on restoration of coastal vegetation in Florida, Environmental Studies Center, Hillsborough Community College, Tampa, FL. p. 13-44.
5. Carlton, Jeffrey M. 1974. Land-building and stabilization by mangroves. Environmental Conservation 1(4):285-294.
6. Carlton, Jeffrey M. and Mark D. Moffler. 1978. Propagation of mangroves by air-layering. Environmental Conservation 5(2):147-150.
7. Carter, Michael R., Lawrence A. Burns, Thomas R. Cavinder, Kenneth R. Dugger, Paul L. Fore, Delbert B. Hicks, H. Lavon Revells, and Thomas W. Schmidt. 1973. Ecosystems analysis of the big cypress swamp and estuaries. U.S. Environmental Protection Agency, Atlanta, GA. EPA 904/9-74-00Z.
8. Chakrabarti K. and A.B. Chaudhuri. 1972. Wildlife biology of the Sunderbans forests: honey production and behaviour pattern of the honeybee. Science and Culture 38(6):269-276.
9. Chandrashekar, M. and M. C. Ball. 1980. Leaf blight of grey mangrove in Australia caused by *Alternaria alternata*. Transactions of the British Mycological Society 75(3):413-418.
10. Chapman, V.J. 1944. 1939 Cambridge University expedition to Jamaica. Journal of the Linnean Society of London 12:407-533.
11. Chapman, V.J. 1970. Mangrove phytosociology. Tropical Ecology 11(1):1-19.
12. Chapman, V.J. 1976. Mangrove vegetation. J. Cramer, Germany. 447 p.
13. Cintron, Gilberto, Ariel E. Lugo, Douglas J. Pool, and Greg Morris. 1978. Mangroves of arid environments in Puerto Rico and adjacent islands. Biotropica 10(2):110-121.
14. Cintron, Gilberto, Ariel E. Lugo, and Ramon Martinez. [In Press]. Structural and functional properties of mangrove forests. In The botany and natural history of Panama. W.G. D'Arcy and M.D.

- Correa A., editors. Missouri Botanical Garden, St. Louis.
15. Clarke, Lesley D. and Nola J. Hannon. 1970. The mangrove swamp and salt marsh communities of the Sydney district III. Plant growth in relation to salinity and waterlogging. *Journal of Ecology* 58: 351-369.
 16. Clough, B.F. and P.M. Attiwill. 1975. Nutrient cycling in a community of *Avicennia marina* in a temperate region of Australia. In: *Proceedings, International Symposium on biology and management of mangroves*. G. Walsh., S.C. Snedaker, H. Teas, editors. University of Florida, Gainesville. p. 137-146.
 17. Davis, John H. 1940. The ecology and geologic role of mangroves in Florida. *Papers of the Tortugas Laboratory (Carnegie Institution)* 32:303-412.
 18. Gill, A. Malcolm. 1971. Endogenous control of growth-ring development in *Avicennia*. *Forest Science* 17(4):462-465.
 19. Halle, F., R.A.A. Oldeman, and P.B. Tomlinson. 1978. *Tropical trees and forests: an architectural analysis*. Springer-Verlag, New York. 441 p.
 20. Hesse, P.R. 1961. Some differences between the soils of *Rhizophora* and *Avicennia* mangrove swamp in Sierra Leone. *Plant and Soil* 14:335-461.
 21. Holdridge, L.H. 1940. Some notes on the mangrove swamps of Puerto Rico. *Caribbean Forester* 1(4): 19-29.
 22. Holdridge, L.H. 1967. *Life zone ecology (revised edition)*. Tropical Science Center, San Jose, Costa Rica. 206 p.
 23. Jimenez, Jorge Arturo. 1981. The mangrove of Costa Rica: a physiognomic characterization. Thesis, MS, University of Miami, Coral Gables, FL. 130 p.
 24. Jenik, Jan. 1978. Roots and root systems in tropical trees: morphologic and ecologic aspects. In *Tropical trees as living systems*. P.B. Tomlinson and Martin H. Zimmermann, editors. Cambridge University Press, Cambridge. p. 323-349.
 25. Karstedt, P., and W. Liese. 1973. Protection of mangrove wood with water-borne preservatives. *Holz als Roh-und Werkstoff* 31(2):73-76.
 26. Lescure, Jean Paul. 1980. Ecological aspects of the mangrove forest in French Guiana. In *Memorias del seminario sobre el estudio científico e impacto humano en el ecosistema de manglares*, UNESCO, Oficina Regional de Ciencia y Tecnología para América Latina y el Caribe. Montevideo, Uruguay. p. 76-93.
 27. Little, E.L. 1961. *Avicennia nitida* (Nomenclature). *Phytologia* 8(2):49-57.
 28. Little, E.L. and F.H. Wadsworth. 1964. *Common trees of Puerto Rico and the Virgin Islands*. U.S. Department of Agriculture, Forest Service, Agriculture Handbook No. 249. Washington, DC. 548 p.
 29. Lugo, Ariel E. 1981. The inland mangroves of Inagua. *Journal of Natural History* 15(5):845-852.
 30. Lugo, Ariel E., and Samuel C. Snedaker. 1974. The ecology of mangroves. *Annual Review of Ecology and Systematics* 5:39-64.
 31. Lugo, Ariel E. and Samuel C. Snedaker. 1975. Properties of a mangrove forest in southern Florida. In *Proceedings, International Symposium on Biological Management of Mangroves*. G. Walsh, S. Snedaker and H. Teas, editors. Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL. p. 170-212.
 32. Lugo, Ariel E., Gary Evink, Mark M. Brinson, Alberto Broce, and Samuel C. Snedaker. 1975. Diurnal rates of photosynthesis, respiration and transpiration in mangrove forests of south Florida. In *Tropical Ecological Systems*. Frank B. Golley and Ernesto Medina, editors. Springer-Verlag, New York, p. 335-350.
 33. Lugo, Ariel E., and Carol Patterson-Zucca. 1977. The impact of low temperature stress on mangrove structure and growth. *Tropical Ecology* 18(2): 149-161.
 34. Lugo, Ariel E., Robert R. Twilley, and Carol Patterson-Zucca. 1980. The role of mangrove forests in the productivity of coastal ecosystems in south Florida. Final report to U.S. Environmental Protection Agency, Corvallis Environmental Research Laboratory, Corvallis, OR. Contract No. R 806079010. Center for Wetlands, University of Florida, Gainesville, FL. 281 p.
 35. Luna Lugo, Anibal. 1976. Manejo de manglares en Venezuela. *Boletín del Instituto Forestal Latino-Americano* 50:41-56.
 36. Marshall, R.C. 1939. *Silviculture of the trees of Trinidad and Tobago*, British West Indies. Oxford University Press London. 247 p.
 37. McMillan, Calvin. 1971. Environmental factors affecting seedling establishment of the black mangrove on the central Texas coast. *Ecology* 52(5):927-929.
 38. Moldenke, Harold. 1973. *Avicenniaceae*. In *Flora of Panama*. Robert E. Woodson, Jr., and Robert W. Schery. *Annals of the Missouri Botanical Garden* 60:149-154.
 39. Moldenke, Harold N. 1975. Additional notes on the genus *Avicennia*. VI. *Phytologia* 32(5):436-457.
 40. Newberry, D. McC. 1980. Infestation of the coccid, *Icerya seychellarum* Westw. on the mangrove *Avicennia marina* (Forsk.) Viernh. on Aldabra Atoll, with special reference to tree age. *Oecologia* 45(3): 325-330.
 41. Pegg, K.G., N.C. Gillespie, and L.I. Forsberg. 1980. *Phytophthora* sp. associated with mangrove death in central coast Queensland. *Australian Plant Pathology* 9(3):6-7.
 42. Piyakarnchana, T. 1981. Severe defoliation of *Avicennia alba* Bl. by larvae of *Cleora injectaria* Walker. *Journal of the Scientific Society of Thailand* 7(1): 33-36.
 43. Puerto Rico Water Resources Authority. 1972. Aguirre power plant complex, environmental report. WRAES-8. 198 p.
 44. Pulver, Terry R. 1976. Transplant techniques for sapling mangrove trees, *Rhizophora mangle*, *Laguncularia racemosa* and *Avicennia germinans* in Florida. *Florida Marine Research Publications* 22:1-14.

45. Rabinowitz, Deborah. 1978. Dispersal properties of mangrove propagules. *Biotropica* 10(1):47-57.
46. Rabinowitz, Deborah. 1978. Mortality and initial propagule size in mangrove seedlings in Panama. *Journal of Ecology* 66:45-51.
47. Record, Samuel J., and Robert W. Hess. 1949. *Timbers of the new world*. Yale University Press, New Haven. 640 p.
48. Rehm, Andrew E. 1976. The effects of the wood-boring isopod *Sphaeroma terebrans* on the mangrove communities of Florida. *Environmental Conservation* 3(1):47-57.
49. Scholander, P.F., L. Van Dam, and S.I. Scholander. 1955. Gas exchange in the roots of mangroves. *American Journal of Botany* 42:92-98.
50. Snedaker, Samuel C., Jorge A. Jimenez, and Melvin S. Brown. 1981. Anomalous aerial roots in *Avicennia germinans* (L.) L. in Florida and Costa Rica. *Bulletin of Marine Science* 31(2):467-470.
51. Soto, Ricardo, and Jorge A. Jimenez. 1982. Analisis fisionomico estructural del manglar de Puerto Soley, La Cruz, Guanacaste, Costa Rica. *Revista Biologica Tropical* 30(2):161-168.
52. Southwell, C.R., and J.D. Bultman. 1971. Marine borer resistance of untreated woods over long periods of immersion in tropical waters. *Biotropica* 3(1):81-107.
53. Surinam Forest Service. 1955. *Surinam Timber*. 2nd edition. Paramaribo. 93 p.
54. Swamy, B.G.L., and D. Padmanabhan. 1961. *Notulae Embryologicae* 1. The functions of endosperm in *Avicennia officinalis*. *Current Science* 11:424-425.
55. Thom, Bruce G. 1967. Mangrove ecology and deltaic geomorphology. Tabasco, Mexico. *Journal of Ecology* 55:301-343.
56. Troup, R.S. 1921. *The silviculture of Indian trees*. Vol. 2. Clarendon Press. Oxford.
57. Truman, R. 1961. The eradication of mangroves. *Aust. J. Sci.* 24:198-199.
58. Twilley, Robert C. 1982. Litter dynamics and organic carbon exchange in black mangrove (*Avicennia germinans*) basin forests in a southwest Florida estuary. PhD Dissertation, University of Florida. 260 p.
59. Uphof, J.C. 1968. *Dictionary of economic plants*. Verlag Von J. Cramer, Germany. 591 p.
60. Wadsworth, F.H., and G.H. Englerth. 1959. Effects of the 1956 hurricane on forests in Puerto Rico. *Caribbean Forester* 20(1 & 2):38-51.
61. Watson, J.G. 1928. Mangrove forests of the Malay Peninsula. *Malay. For. Rec.* 6(24):125-149.
62. Weaver, Peter L. 1979. Tree growth in several tropical forests of Puerto Rico. U.S. Dep. Agric. For. Ser. Res. Pap. SO-152. South. For. Exp. Stn. New Orleans, La. 15 p.
63. Wells, John T., and James M. Coleman. 1981. Periodic mudflat progradation, northeastern coast of South America: a hypothesis. *Jour. Sed. Petrol.* 51(4):1069-1075.
64. West, Robert C. 1977. Tidal salt-marsh and mangal formations of Middle and South America. In *Ecosystems of the world. 1 Wet coastal ecosystems*. V.J. Chapman, ed. Elsevier Scientific Publishing Co., Oxford. p. 193-213.